



STUDIO PRELIMINARE DI PROCESSI
 $\beta\beta$ IN ^{106}Cd USANDO UN CRISTALLO
SCINTILLATORE ARRICCHITO DI $^{106}\text{CdWO}_4$
IN COINCIDENZA/ANTICOINCIDENZA CON
DUE CdWO_4

107° Congresso Nazionale della Società Italiana di Fisica
13-17 Settembre 2021

Alice Leoncini
alice.leoncini@roma1.infn.it
INFN-ROMA


Introduction to double beta decay

- It can be studied in even-even nuclei when the single β decay is energetically forbidden due to the pairing interaction.

- For example: $^{106}_{48}\text{Cd} \rightarrow ^{106}_{46}\text{Pd} + 2e^+ + 2\nu_e$

$$2\nu 2\beta^+ : \frac{A}{Z}X \rightarrow \frac{A}{Z-2}X + 2e^+ + 2\nu_e \quad \text{L conserved}$$

$$0\nu 2\beta^+ : \frac{A}{Z}X \rightarrow \frac{A}{Z-2}X + 2e^+ \quad \text{L violated } (\Delta L = 2)$$



- If the total lepton number L is violated, also the following $\beta^+\beta^+$ processes are possible:

$$0\nu\epsilon\beta^+ : e^- + \frac{A}{Z}X \rightarrow \frac{A}{Z-2}X + e^+ + \text{X-rays}$$


$$0\nu 2\epsilon : e^- + e^- + \frac{A}{Z}X \rightarrow \frac{A}{Z-2}X^* \rightarrow \frac{A}{Z-2}X + \gamma + \text{X-rays}$$

- Easier to identify: Positron(s) annihilation gives rise to 511 keV γ rays.
- $\epsilon\beta^+$ and $2\beta^+$ can clarify the possible contribution of the right-handed currents to the $0\nu\beta^-\beta^-$ decay rate;
- Possibility of a resonant $0\nu 2\epsilon$ process \rightarrow in case of close degeneracy of the initial and final (excited) nuclear states $\rightarrow \frac{1}{T_{1/2}} \propto \frac{\Gamma}{(Q-E)^2 + \Gamma^2/4}$


DOUBLE BETA DECAY IN ^{106}Cd

Advantages in the use of ^{106}Cd :


- One of the biggest decay energy: $Q_{\beta\beta} = (2775.39 \pm 0.10)$ keV;
- High isotopic abundance: $\delta = (1.245 \pm 0.022)$ %;
- Favorable theoretical predictions for half-lives for some 2ν modes ($T_{1/2} \sim 10^{21} - 10^{22}$ yr) that could be reached by modern low-counting techniques;
- Possibility of resonant 2ϵ to excited levels of ^{106}Pd ;
- Possibility of enrichment by gas centrifugation, existing technologies of cadmium purification and availability of Cd-containing detectors to realize calorimetric experiments with a high detection efficiency.



SEARCHES FOR $\beta\beta$ DECAY IN ^{106}Cd at GRAN SASSO: PREVIOUS STAGES OF THE EXPERIMENT




A $^{106}\text{CdWO}_4$ crystal (215.4 g) enriched in ^{106}Cd at 66 % was grown and used in three previous stages of the experiment:




Stage 1 (2012): $^{106}\text{CdWO}_4$ crystal was fixed inside a cavity in the central part of a polystyrene light-guide. The experimental apparatus was located in the DAMA/R&D setup at LNGS.

Stage 2 (2016): $^{106}\text{CdWO}_4$ crystal in coincidence with 4 ultra-low-background HPGe detectors of the GeMulti setup of the STELLA (SubTERRanean Low Level Assay) facility at LNGS.




Stage 3 (2020): at low background DAMA/CRYSTAL setup located at LNGS. $^{106}\text{CdWO}_4$ detector in coincidence with two large-volume CdWO_4 scintillators detectors in close geometry to improve the detection efficiency to γ quanta emitted in the $\beta\beta$ processes in ^{106}Cd .





THE NEW EXPERIMENT INSTALLED IN THE DAMA/R&D SETUP AT LNGS

- $^{106}\text{CdWO}_4$ is housed in a cylindrical cut-out of the two CdWO_4 scintillators which almost completely envelop the enriched crystal.
- An event-by-event DAQ records pulses in case of:
 - an event with $E > 500$ keV in $^{106}\text{CdWO}_4$ detector;
 - $^{106}\text{CdWO}_4$ detector in coincidence with at least one of the CdWO_4 counters.



Monte Carlo simulation: Radioactive Contaminants

❖ **^{238}U chain:** $^{238}\text{U} \rightarrow ^{234}\text{U}$, $^{234}\text{U} \rightarrow ^{230}\text{Th}$, $^{230}\text{Th} \rightarrow ^{226}\text{Ra}$, $^{226}\text{Ra} \rightarrow ^{210}\text{Pb}$, $^{210}\text{Pb} \rightarrow ^{206}\text{Pb}$.

❖ **^{232}Th chain:** $^{232}\text{Th} \rightarrow ^{228}\text{Ra}$, $^{228}\text{Ra} \rightarrow ^{228}\text{Th}$, $^{228}\text{Th} \rightarrow ^{208}\text{Pb}$.


❖ ^{40}K

These were simulated in the following materials of the setup:

- 1) $^{106}\text{CdWO}_4$;
- 2) the two natural CdWO_4 crystals;
- 3) Plastic light-guide;
- 4) the optical couplants;
- 5) the teflon tapes;
- 6) the teflon details, which include the teflon spring and support 1 and 2;


- 7) the two quartz light-guides connected to CdWO_4 s;
- 8) the quartz light-guide connected to $^{106}\text{CdWO}_4$;
- 9) the "copper internal" volume;
- 10) the "copper external" volume;
- 11) the PMTs coupled to CdWO_4 s;
- 12) the PMT coupled to the $^{106}\text{CdWO}_4$ detector.

Also: ^{56}Co , ^{60}Co in copper internal and ^{113}Cd , $^{113\text{m}}\text{Cd}$ in the three CdWO_4 crystals.



Data taking and energy calibration



- The energy scale was calibrated using ^{22}Na , ^{60}Co , ^{133}Ba , ^{137}Cs , and ^{228}Th γ sources. The data taking started in October 2019 and it is still in progress. In this work, a total time of accumulated data of 466.64 d is considered.




CdWO_4 -1


$$\sigma_\gamma = p_1 \sqrt{E_\gamma} \text{ where } E_\gamma \text{ is expressed in keV.}$$

CdWO_4 -2



$^{106}\text{CdWO}_4$





PULSE SHAPE DISCRIMINATION OF α AND $\gamma(\beta)$ EVENTS

- The difference in CdWO_4 scintillation pulse shape for β particles (γ quanta) and α particles can be used in order to suppress the background caused by α radioactive contamination of the detectors due to the residual contamination in ^{232}Th and ^{238}U with their daughters.

$$SI = \sum f(t_k) \times P(t_k) / \sum f(t_k)$$

The mean value of the shape indicator vs energy is represented together with 1σ intervals for the $^{106}\text{CdWO}_4$ detector and also 2σ , 3σ intervals for the CdWO_4 scintillators.

EXPERIMENTAL SPECTRA AFTER EVENT SELECTIONS

❖ Anticoincidence mode (AC):


- E (both CdWO_4) < 100 keV
- $SI > \mu_{SI\beta} - 1\sigma_{SI\beta}$ for the $^{106}\text{CdWO}_4$ detector
- PSD efficiency: $\eta_\beta = 0.84$

❖ Coincidence mode (CC511):



- E (one of the two CdWO_4) $= (511 \pm 2\sigma)$ keV
- $SI > \mu_{SI\beta} - 3\sigma_{SI\beta}$ for the two CdWO_4 crystals
- NO PSD efficiency

❖ Double coincidence mode (CC511&511):

- E (both CdWO_4) $= (511 \pm 2\sigma)$ keV
- $SI > \mu_{SI\beta} - 3\sigma_{SI\beta}$ for the two CdWO_4 crystals
- NO PSD efficiency



Energy spectra
of $\beta(\gamma)$ events





ENERGY SPECTRA WITH BACKGROUND MODEL

- The χ^2 function used for the fit was estimated with the maximum likelihood estimator, which takes into account the Poissonian nature of the fluctuations in the experimental bins.
- For each contaminant considered, the relative model was introduced for each of the two spectra, but with a **single free parameter** (proportional to the activity of the contaminant).

Radioactive contamination (mBq/kg) of the materials of the setup.

Material	²³⁸ U	²²⁶ Ra	²²⁸ Ac	²²⁸ Th	⁴⁰ K
¹⁰⁶ CdWO ₄	0.60(2)	0.018(8)	-	0.034(12)	<1.6
CdWO ₄ crystals	-	-	-	-	<0.5
Plastic light-guide	-	<20	<30	<5	<18
Quartz light-guides	-	<6	<15	<1.2	<6
Copper	-	<0.4	<1.0	<0.10	<0.5
PMTs	-	<1000	<4000	<160	<1600



Search for 2ϵ , $\epsilon\beta^+$ and $2\beta^+$ processes in ^{106}Cd

- The measured energy spectrum does not contain peculiarities which could be ascribed to $\beta\beta$ decay processes in ^{106}Cd . Therefore, the data have been analyzed estimating lower half-life limits, using the following formula:

$$\lim T_{1/2} = N \cdot \eta \cdot t \cdot \ln 2 / \lim S$$

where:

- N is the number of ^{106}Cd nuclei in the $^{106}\text{CdWO}_4$ crystal ($N = 2.42 \times 10^{23}$);
- η is the detection efficiency for the process of decay (calculated as a ratio of the events number in the signal model which satisfies the investigated experimental condition, to the number of generated events);
- t is the time of measurements (466.64 d);
- $\lim S$ is the number of events of the effect searched for, which can be excluded at a given confidence level (C.L.; all limits on $\beta\beta$ processes in ^{106}Cd are given at the 90% C.L. in the present study).

SEARCH FOR $\beta\beta$ PROCESSES THROUGH THE STUDY OF TRIPLE COINCIDENCES

- Spectrum of $^{106}\text{CdWO}_4$ in coincidence with events at energy $E = (511 \pm 2\sigma)$ keV in both of the two natural CdWO_4 scintillators → **NO events in the energy region >520 keV** → **lim $S = 2.3$ (90% C.L.)**

Decay, Level of ^{106}Pd	η	limit $T_{1/2}$ [yr] at 90% C.L.
$0\nu 2\epsilon$ 2^+512	0.003	$\geq 2.6 \times 10^{20}$
$2\nu\epsilon\beta^+$ g.s.	0.016	$\geq 1.6 \times 10^{21}$
$2\nu\epsilon\beta^+$ 2^+512	0.037	$\geq 3.5 \times 10^{21}$
$2\nu\epsilon\beta^+$ 2^+1128	0.023	$\geq 2.1 \times 10^{21}$
$2\nu\epsilon\beta^+$ 0^+1134	0.027	$\geq 2.6 \times 10^{21}$
$0\nu\epsilon\beta^+$ g.s.	0.035	$\geq 3.3 \times 10^{21}$
$0\nu\epsilon\beta^+$ 2^+512	0.043	$\geq 4.0 \times 10^{21}$
$0\nu\epsilon\beta^+$ 2^+1128	0.028	$\geq 2.6 \times 10^{21}$
$0\nu\epsilon\beta^+$ 0^+1134	0.030	$\geq 2.8 \times 10^{21}$
$2\nu 2\beta^+$ g.s.	0.050	$\geq 4.6 \times 10^{21}$
$2\nu 2\beta^+$ 2^+512	0.046	$\geq 4.3 \times 10^{21}$
$0\nu 2\beta^+$ g.s.	0.051	$\geq 4.7 \times 10^{21}$
$0\nu 2\beta^+$ 2^+512	0.046	$\geq 4.3 \times 10^{21}$

SEARCH FOR $\beta\beta$ PROCESSES BY FITTING THE MEASURED SPECTRA

- Simulated $\beta\beta$ models were added to the background model in the fit of the AC spectrum plus the CC511 spectrum.
- $\beta\beta$ decay model summed up to the background model is normalized to one decay, i.e. it is divided by the total number of generated decays (and the PSD cut efficiency is also taken into account).
- Therefore the fit directly returns the total number of decays attributable to the searched process i.e. S/η .
- According to Feldman & Cousins[127] procedure, an upper limit on S/η (90% C.L.) is calculated.

Decay, Level of ^{106}Pd	S/η	$lim(S/\eta)$	limit $T_{1/2}$ [yr] at 90% C.L.
$0\nu 2\epsilon$ g.s.	194 ± 105	367	$\geq 2.5 \times 10^{20}$
Res. $0\nu 2K$ 2718	-22 ± 143	214	$\geq 4.3 \times 10^{20}$
Res. $0\nu KL_1$ 4^+2741	132 ± 102	300	$\geq 3.1 \times 10^{20}$
Res. $0\nu KL_3$ 2, 3-2748	-39 ± 98	124	$\geq 7.4 \times 10^{20}$

[127] Feldman, G.J.; Cousins, R.D. Unified approach to the classical statistical analysis of small signals. *Phys. Rev. D* 57, 1998, 3873-3889.

CONCLUSIONS & RESULTS

Decay, Level of ^{106}Pd	Exp. selection	limit $T_{1/2}$ [yr] at 90% C.L.	
		Best present	Best previous
$0\nu 2\epsilon$ g.s.	AC&CC511	$\geq 2.5 \times 10^{20}$	$\geq 1.0 \times 10^{21}$ [10]
$0\nu 2\epsilon$ 2^+512	CC511&511	$\geq 2.6 \times 10^{20}$	$\geq 5.1 \times 10^{20}$ [10]
Res. $0\nu 2K$ 2718	AC&CC511	$\geq 4.3 \times 10^{20}$	$\geq 2.9 \times 10^{21}$ [12]
Res. $0\nu KL_1$ 4^+2741	AC&CC511	$\geq 3.1 \times 10^{20}$	$\geq 9.0 \times 10^{20}$ [10]
Res. $0\nu KL_3$ $2,3^-2748$	AC&CC511	$\geq 7.4 \times 10^{20}$	$\geq 1.4 \times 10^{21}$ [11]
$2\nu\epsilon\beta^+$ g.s.	CC 511&511	$\geq 1.6 \times 10^{21}$	$\geq 2.1 \times 10^{21}$ [12]
$2\nu\epsilon\beta^+$ 2^+512	CC 511&511	$\geq 3.5 \times 10^{21}$	$\geq 2.7 \times 10^{21}$ [12]
$2\nu\epsilon\beta^+$ 2^+1128	CC 511&511	$\geq 2.1 \times 10^{21}$	$\geq 1.3 \times 10^{21}$ [12]
$2\nu\epsilon\beta^+$ 0^+1134	CC 511&511	$\geq 2.6 \times 10^{21}$	$\geq 1.1 \times 10^{21}$ [11]
$0\nu\epsilon\beta^+$ g.s.	CC511&511	$\geq 3.3 \times 10^{21}$	$\geq 1.4 \times 10^{22}$ [12]
$0\nu\epsilon\beta^+$ 2^+512	CC511&511	$\geq 4.0 \times 10^{21}$	$\geq 9.7 \times 10^{21}$ [12]
$0\nu\epsilon\beta^+$ 2^+1128	CC511&511	$\geq 2.6 \times 10^{21}$	$\geq 1.1 \times 10^{22}$ [12]
$0\nu\epsilon\beta^+$ 0^+1134	CC 511&511	$\geq 2.8 \times 10^{21}$	$\geq 1.9 \times 10^{21}$ [11]
$2\nu 2\beta^+$ g.s.	CC 511&511	$\geq 4.6 \times 10^{21}$	$\geq 2.3 \times 10^{21}$ [11]
$2\nu 2\beta^+$ 2^+512	CC 511&511	$\geq 4.3 \times 10^{21}$	$\geq 2.5 \times 10^{21}$ [11]
$0\nu 2\beta^+$ g.s.	CC511&511	$\geq 4.7 \times 10^{21}$	$\geq 5.9 \times 10^{21}$ [12]
$0\nu 2\beta^+$ 2^+512	CC511&511	$\geq 4.3 \times 10^{21}$	$\geq 4.0 \times 10^{21}$ [12]

[10] Belli, P.; Bernabei, R.; Boiko, R.S.; Brudanin, V.B.; Cappella, F.; Caracciolo, V.; Cerulli, R.; Chernyak, D.M.; Danevich, F.A.; d'Angelo, S. *et al.* Search for double- β decay processes in ^{106}Cd with the help of a $^{106}\text{CdWO}_4$ crystal scintillator. *Phys. Rev. C* 85, 2012, 044610.

[11] Belli, P.; Bernabei, R.; Brudanin, V.B.; Cappella, F.; Caracciolo, V.; Cerulli, R.; Chernyak, D.M.; Danevich, F.A.; d'Angelo, S.; Di Marco, A.; *et al.* Search for double- β decay in ^{106}Cd with an enriched $^{106}\text{CdWO}_4$ crystal scintillator in coincidence with four HPGe detectors. *Phys. Rev. C* 93, 2016, 045502.

[12] Belli, P. *et al.* Search for Double Beta Decay of ^{106}Cd with an Enriched $^{106}\text{CdWO}_4$ Crystal Scintillator in Coincidence with CdWO_4 Scintillation Counters, *Universe* 6, 2020, 182.

- The highest sensitivity to several decay channels with positron(s) emission was achieved using the data that were gathered by the $^{106}\text{CdWO}_4$ detector in coincidence with 511 keV annihilation γ quanta in both of the two CdWO_4 counters.
- Limits have been improved to a factor 2-3 with respect to the previous experiments.
- The sensitivity obtained on the $T_{1/2}$ for the case $2\nu\epsilon\beta^+$ approaches the theoretical predictions: $T_{1/2} \sim 10^{21} - 10^{22}$ yr.
- The experiment is still running with the purpose of improving the sensitivity to all the decay channels of ^{106}Cd .