

RECENT DEVELOPMENTS AND RESULTS ON $\beta\beta$ DECAYS WITH CRYSTAL SCINTILLATORS AND HP-GE SPECTROMETRY



**“IS QUANTUM THEORY EXACT? THE QUEST FOR THE SPIN-
STATISTICS CONNECTION VIOLATION AND RELATED ITEMS”**


FRASCATI 2-5.07.2018

**Dr. A. Di Marco
INFN Roma “Tor Vergata”**



DAMA SET-UPS

an observatory for rare processes @ LNGS



Collaboration:

Roma Tor Vergata, Roma La Sapienza, LNGS, IHEP/Beijing

+ by-products and small scale expts.: INR-Kiev + other institutions

+ neutron meas.: ENEA-Frascati, ENEA-Casaccia


+ in some studies on $\beta\beta$ decays (DST-MAE & Inter-Univ. project): IIT Kharagpur and Ropar, India

web site: <http://people.roma2.infn.it/dama>

MAIN RESULTS OBTAINED BY DAMA IN THE SEARCH FOR RARE PROCESSES

- First or improved results in the search for 2β decays of ~ 30 candidate isotopes: ^{40}Ca , ^{46}Ca , ^{48}Ca , ^{64}Zn , ^{70}Zn , ^{100}Mo , ^{96}Ru , ^{104}Ru , ^{106}Cd , ^{108}Cd , ^{114}Cd , ^{116}Cd , ^{112}Sn , ^{124}Sn , ^{134}Xe , ^{136}Xe , ^{130}Ba , ^{136}Ce , ^{138}Ce , ^{142}Ce , ^{156}Dy , ^{158}Dy , ^{180}W , ^{186}W , ^{184}Os , ^{192}Os , ^{190}Pt and ^{198}Pt (observed $2\nu 2\beta$ decay in ^{100}Mo , ^{116}Cd)
- The best experimental sensitivities in the field for 2β decays with positron emission (^{106}Cd)

First observation of α decays of ^{151}Eu with a $\text{CaF}_2(\text{Eu})$ scintillator and of ^{190}Pt to the first excited level ($E_{\text{exc}}=137.2$ keV) of ^{186}Os



$(T_{1/2}=5 \times 10^{18}\text{yr})$

Observation of correlated e^+e^- pairs emission in α decay of ^{241}Am ($A_{e^+e^-}/A_\alpha \approx 5 \times 10^{-9}$)

Search for cluster decays of ^{127}I , ^{138}La and ^{139}La


Search for PEP violating processes in Sodium and in Iodine

Search for N, NN, NNN decay into invisible channels in ^{129}Xe and ^{136}Xe

CNC processes, e.g. in ^{127}I , ^{136}Xe , ^{100}Mo and ^{139}La

Search for ^7Li solar axions using resonant absorption in LiF crystal

Investigations of rare β decays of ^{113}Cd ($T_{1/2}=8 \times 10^{15}\text{yr}$), $^{113\text{m}}\text{Cd}$ with CdWO_4 scintillator and ^{48}Ca with a $\text{CaF}_2(\text{Eu})$ detector



Search for spontaneous transition of ^{23}Na and ^{127}I nuclei to superdense state


Dark Matter investigation

... many others are in progress

SOME EXAMPLE OF ISOTOPES DECAYING $\beta\beta$ THAT COULD BE STUDIED BY CRYSTAL SCINTILLATORS WITH «SOURCE=DETECTOR» APPROACH

Isotops	Nat.Ab.(%)	Q(keV)	Decay Mode	Scintillator
^{64}Zn	8.63	1095.7	$\varepsilon\beta^+, 2\varepsilon$	ZnWO_4
^{70}Zn	0.62	998.5	$2\beta^-$	ZnWO_4
^{180}W	0.12	144	2ε	$\text{ZnWO}_4, \text{CdWO}_4, \text{PbWO}_4$
^{186}W	28.43	489.9	$2\beta^-$	$\text{ZnWO}_4, \text{CdWO}_4, \text{PbWO}_4$
^{106}Cd	1.25	2771	$2\beta^+, \varepsilon\beta^+, 2\varepsilon$	$^{106}\text{CdWO}_4$
^{108}Cd	0.89	269	2ε	CdWO_4
^{114}Cd	28.73	536.8	$2\beta^-$	CdWO_4
^{116}Cd	7.49	2805	$2\beta^-$	$^{116}\text{CdWO}_4$
^{40}Ca	96.941	193.78	2ε	$\text{CaF}_2, \text{CaMoO}_4$
^{46}Ca	0.004	990.4	$2\beta^-$	$\text{CaF}_2, \text{CaMoO}_4$
^{48}Ca	0.187	4272	$2\beta^-$	$\text{CaF}_2, \text{CaMoO}_4$
^{136}Ce	0.185	2419	$2\beta^+, \varepsilon\beta^+$	$\text{CeCl}_3, \text{CeF}_3, \text{CeBr}_3$
^{138}Ce	0.251	693	2ε	$\text{CeCl}_3, \text{CeF}_3, \text{CeBr}_3$
^{142}Ce	11.114	1416.9	$2\beta^-$	$\text{CeCl}_3, \text{CeF}_3, \text{CeBr}_3$
^{130}Ba	0.106	2611	$2\beta^+, \varepsilon\beta^+, 2\varepsilon$	$\text{BaF}_2, \text{BaCl}_2(\text{Eu}), \text{BaI}_2(\text{Eu})$
^{92}Mo	14.84	1649	$\varepsilon\beta^+, 2\varepsilon$	$\text{PbMoO}_4, \text{LiMoO}_4, \text{CaMoO}_4$
^{100}Mo	9.63	3034	$2\beta^-$	$\text{PbMoO}_4, \text{LiMoO}_4, \text{CaMoO}_4$
^{84}Sr	0.56	1786.8	$\varepsilon\beta^+$	$\text{SrCl}_2, \text{SrI}_2(\text{Eu})$

DAMA/R&D + DAMA/CRYS - SOME MAIN PREVIOUS RESULTS:



- scintillators developments: radio-purification enrichment, optical features, etc.
- exploiting the potentiality of the low background scintillation technique to investigate rare processes with high sensitivity
- realization of pilot experiments

Some example:

$\text{CaF}_2(\text{Eu})$, CeF_3 , BaF_2 , CdWO_4 , $^{106}\text{CdWO}_4$, $^{116}\text{CdWO}_4$,
 ZnWO_4 , $\text{LaCl}_3(\text{Ce})$, $\text{LiEu}(\text{BO}_3)_3$, $\text{LiF}(\text{W})$, CeCl_3 , Li_2MoO_4 ,
 $\text{SrI}_2(\text{Eu})$, etc.



AP7(1997)73, N.Cim.A110(1997)189, NPB563(1999)97,
AP10(1999)115, NPA705(2002)29, NIMA498(2003)352,
NIMA525(2004)535, NIMA555(2005)270, UJP51(2006)1037,
NPA789(2007)15, PRC76(2007)064603, PLB658(2008)193,
EPJA36(2008)167, NPA824(2009)101, NPA826(2009)256,
JPG:NPP38(2011)115107, JPG: NPP38(2011)015103,
JINST6(2011)P08011, PRC85(2012)044610, EPJC73(2013)2276,
EPJA50(2014)134, PS90(2015)085301

DAMA/GE AND LNGS STELLA FACILITY

Ge detectors used by DAMA in previous searches:

DAMA/Ge (GeBer)

- 244 cm³ n-type HPGe detector
- Thin Carbon window: 0.76 mm thickness

GeCris

- 465 cm³ p-type HPGe detector
- Thin Cu window: 1 mm thickness

GeMulti

- Four 225 cm³ p-type HPGe detectors mounted in one cryostat with a well in the center
- Thin Al window: 1.3 mm thickness

GeBEGe

- Broad Energy Ge detector (especially designed for low energy γ spectrometry)
- Thin Cu window: 1.5 mm thickness

Typical shield from environmental radioactivity

- 5-10 cm of OFHC copper
- 5 cm of low activity lead (< 3 Bq/kg of ²¹⁰Pb)
- 15-25 cm of lead
- 10 cm of borated polyethylene (GeBer)
- Air-tight PMMA box flushed with HP nitrogen



DAMA results

- Search for $\beta\beta$ decays of many candidate isotopes (next slide)
- Search for ⁷Li solar axions ([NPA806\(2008\)388](#), [PLB711\(2012\)41](#))
- First observation of α decay of ¹⁹⁰Pt to the first excited level of ¹⁸⁶Os ([PRC83\(2011\)034603](#))
- Qualification of many materials: e.g. CdWO₄, ZnWO₄ ([NIMA626-7\(2011\)31](#), [NIMA615\(2010\)301](#)), Li₆Eu(BO₃)₃ ([NIMA572\(2007\)734](#)), Li₂MoO₄ ([NIMA607\(2009\)573](#)), SrI₂(Eu) ([NIMA670\(2012\)10](#)), ⁷LiI(Eu) ([NIMA704\(2013\)40](#))

First or improved results for 2β decays of many isotopes

^{136}Ce $Q_{\beta\beta}=2378.55$ keV; 2ε , $\varepsilon\beta^+$, $2\beta^+$; ^{138}Ce $Q_{\beta\beta}=691$ keV; 2ε

- CeO_2 sample (627 g) in GeCris detector (2299 h) $\Rightarrow T_{1/2}$ limits: 10^{17} - 10^{19} yr [Eur. Phys. J. A 53 (2017) 172]
- CeO_2 sample (732 g) in GeCris detector (1900 h) $\Rightarrow T_{1/2}$ limits: 10^{17} - 10^{18} yr [Nucl. Phys. A 930 (2014) 195]
- CeCl_3 crystal (6.9 g) in DAMA/Ge detec. (1280 h) $\Rightarrow T_{1/2}$ limits: $(1\div 6)10^{15}$ yr [Nucl. Phys. A 824 (2009) 101]

^{106}Cd $Q_{\beta\beta}=2775.39$ keV; 2ε (res 0ν), $\varepsilon\beta^+$, $2\beta^+$ [Phys. Rev. C 93 (2016) 045502]

- $^{106}\text{CdWO}_4$ crystal scintillator (216 g) in GeMulti (13085 h) $\Rightarrow T_{1/2}$ limits: 10^{20} - 10^{21} yr

^{96}Ru $Q_{\beta\beta}=2714.51$ keV; 2ε (res 0ν), $\varepsilon\beta^+$, $2\beta^+$, ^{104}Ru $Q_{\beta\beta}=1301.2$ keV; $2\beta^-$

- Purified Ru samples in GeMulti det. (0.56kg \times yr) $\Rightarrow T_{1/2}$ limits: 10^{20} - 10^{21} yr [Phys. Rev. C 87 (2013) 034607]
- Ru sample (473 g) in GeCrys detector (158 h) $\Rightarrow T_{1/2}$ limits: 10^{18} - 10^{19} yr [Eur. Phys. J. A 42 (2009) 171]

^{184}Os $Q_{\beta\beta}=1453.7$ keV; 2ε (res 0ν), $\varepsilon\beta^+$; ^{192}Os $Q_{\beta\beta}=412.4$ keV; $2\beta^-$ [Eur. Phys. J. A 49 (2013) 24]

- Os sample (173 g) in GeCris detector (2741 h) $\Rightarrow T_{1/2}$ limits: 10^{16} - 10^{17} yr for ^{184}Os and 10^{19} yr for ^{192}Os

^{190}Pt $Q_{\beta\beta}=1383$ keV; 2ε (res 0ν), $\varepsilon\beta^+$; ^{198}Pt $Q_{\beta\beta}=1049$ keV; $2\beta^-$ [Eur. Phys. J. A 47 (2011) 91]

- Pt sample (42.5 g) in GeCris detector (1815 h) $\Rightarrow T_{1/2}$ limits: 10^{14} - 10^{16} yr for ^{190}Pt and 10^{18} yr for ^{198}Pt

^{156}Dy $Q_{\beta\beta}=2005.95$ keV; 2ε , $\varepsilon\beta^+$; ^{158}Dy $Q_{\beta\beta}=282.7$ keV; 2ε [Nucl. Phys. A 859 (2011) 126]



- Dy_2O_3 sample (322 g) in DAMA/Ge det. (2512 h) $\Rightarrow T_{1/2}$ limits: 10^{14} - 10^{16} yr

^{100}Mo $Q_{\beta\beta}=3035$ keV; $2\beta^-$ [Nucl. Phys. A 846 (2010) 143]

- $^{100}\text{MoO}_3$ sample (1199 g) enriched in ^{100}Mo at 99.5% in GeMulti detector
 \Rightarrow observation of $^{100}\text{Mo} \rightarrow ^{100}\text{Ru}(0_1^+)$ decay: $T_{1/2} = 6.9_{-0.8}^{+1.0}(\text{stat}) \pm 0.7(\text{syst}) \times 10^{20}$ yr

The best experimental sensitivities in the field for 2β decays with positron emission

Summary of searches for $\beta\beta$ decay modes in various isotopes (partial list)



ARMONIA: New observation of $2\nu 2\beta^-$ $^{100}\text{Mo} \rightarrow ^{100}\text{Ru}$ (g.s. $\rightarrow 0_1^+$) decay NPA846 (2010)143

AURORA: New observation of $2\nu 2\beta^-$ ^{116}Cd decay J.Phys.:Conf.Ser.718(2016)062009

- Many competitive limits obtained on lifetime of $2\beta^+$, $\epsilon\beta^+$ and 2ϵ processes (^{40}Ca , ^{64}Zn , ^{96}Ru , ^{106}Cd , ^{108}Cd , ^{130}Ba , ^{136}Ce , ^{138}Ce , ^{180}W , ^{190}Pt , ^{184}Os , ^{156}Dy , ^{158}Dy , ...).
- First searches for resonant $0\nu 2\epsilon$ decays in some isotopes



Armonia


NPA846(2010)143

(meAsuReMent of twO-NeutrIno $\beta\beta$ decAy of ^{100}Mo to 0^+_1 level of ^{100}Ru)

In addition to the transition to the g.s., the $2\beta 2\nu$ decay of ^{100}Mo was registered also for the transition to the first excited 0^+_1 level of ^{100}Ru

If 0^+_1 excited level of ^{100}Ru ($E=1130$ keV) populated
 \Rightarrow two γ quanta (591 keV + 540 keV) emitted in cascade


$^{100}\text{MoO}_3$ sample (mass = 1199 g) enriched in ^{100}Mo at 99.5% installed in GeMulti setup



$T_{1/2}$ measured in several experiments:

	$T_{1/2}$, 10^{20} yr	Year [Ref.]
Frejus UL (4800 m w.e.), HP Ge 100 cm^3 , 994 g of ^{100}Mo (99.5%), 2298 h, only 1-d spectrum;	> 12	1992 [19]
Soudan mine (2090 m w.e.), HP Ge 114 cm^3 , 956 g of ^{100}Mo (98.5%), 9970 h, 1-d spectrum;	$6.1^{+1.8}_{-1.1}$	1995 [11] ^a
Modane UL (4800 m w.e.), 4 HP Ge detectors (100, 120, 380, 400 cm^3), 17 different ^{100}Mo samples (107–1005 g, 95.1–99.3%, 142–1599 h), sum of 1-d spectra;	$9.3^{+2.8}_{-1.7}$	1999 [14]
Modane UL (4800 m w.e.), NEMO-3 detector, 6914 g of ^{100}Mo foils in 12 sectors (95.1–98.9%), 8024 h, individual energies of γ and e^- , tracks for e^- ;	$5.7^{+1.5}_{-1.2}$	2007 [15]
Ground level (10 m w.e.), 2 HP Ge detectors (300 cm^3) in coincidence, 1050 g of ^{100}Mo (98.4%), 21720 h, coincidence spectrum;	$5.5^{+1.2}_{-0.9}$	2009 [16] ^b
Gran Sasso UL (3600 m w.e.), 4 HP Ge detectors (225 cm^3 each) in coincidence, 1199 g of $^{100}\text{MoO}_3$ (99.5%), 18120 h, coincidence and 1-d spectra.	$6.9^{+1.2}_{-1.1}$	This work

Aim of the experiment: remeasurement of the Mo sample used before in the Frejus exp. (not in agreement with other results)



Armonia Results

NPA846(2010)143

Sum spectrum of all 4 HPGe detectors

1-dimensional energy spectrum analysis

Both peaks at 540 keV and 591 keV expected for $2\beta 2\nu$ decay $^{100}\text{Mo} \rightarrow ^{100}\text{Ru}(0_1^+)$ are observed in the data collected with $^{100}\text{MoO}_3$


In the background spectrum they are absent

Fit of peak @ 539.5 keV: $E = 539.4 \pm 0.2$ keV; $S_{540} = 319 \pm 56$ events

Fit of peak @ 590.8 keV: $E = 590.9 \pm 0.2$ keV; $S_{591} = 278 \pm 53$ events

$$T_{1/2} = 6.9_{-0.8}^{+1.0}(\text{stat.}) \pm 0.7(\text{syst.}) \times 10^{20} \text{ yr.}$$

Most of systematic unc. due to calculation of the efficiencies



2-dim energy spectrum analysis

Double coincidences when fixing the energy of one of the Ge detectors

Eight events detected (red)


$$T_{1/2} = 6.8_{-1.8}^{+3.7}(\text{stat.}) \times 10^{20} \text{ yr}$$

in agreement with the half life derived in 1-d analysis

EI = (540±2) keV

EI = (591±2) keV

EI = (545±2) keV
(background)



Investigation of 2β decay of ^{116}Cd with enriched $^{116}\text{CdWO}_4$ crystal scintillators

^{116}Cd

One of the best isotope for $0\nu 2\beta$ decay search:


- $Q_{\beta\beta} = 2813.44(13)$ keV
- $\delta = 7.49(18)\%$
- possible high isotopic enrichment
- promising theoretical calculation

$^{116}\text{CdWO}_4$ crystal scintillators

Grown by the low-thermal-gradient Czochralski technique after deep purification of ^{116}Cd and W;


+ annealing to improve the optical transmission curve

- ✓ Good optical and scintillation properties
- ✓ $^{116}\text{CdWO}_4$ crystals **enriched at 82%**
- ✓ Active source approach (high detection efficiency)
- ✓ Low levels of internal contamination in (U,Th, K)
- ✓ α/β discrimination capability



Recrystalliz.




AURORA exp.



THE AURORA EXPERIMENT IN THE DAMA/R&D SET-UP


Two enriched $^{116}\text{CdWO}_4$ crystal scintillators
 (total mass: 1.162 kg, ^{116}Cd @ 82%)

- ✓ Started in 2011
- ✓ Upgrade - March 2014
- ✓ Total live time since 2014: 25037 h
- ✓ Background level at 2.7-2.9 MeV: **0.1 counts/keV/kg/yr**



BACKGROUND IDENTIFICATION 1: PULSE SHAPE DISCRIMINATION

Event-by-event DAQ based on a 1 GS/s 8 bit transient digitizer (operated at 50 MS/s) records the pulse shape over a time window of 100 μ s from the $^{116}\text{CdWO}_4$ detectors



T=25037 h
M=1.162 kg


$$SI = \sum f(t_k) \times P(t_k) / \sum f(t_k)$$

$$P(t) = [f_\alpha(t) - f_\gamma(t)] / [f_\alpha(t) + f_\gamma(t)]$$

$f(t_k)$ \rightarrow amplitude at t_k

$P(t_k)$ \rightarrow weight function

$f_{\alpha,\gamma}(t_k)$ \rightarrow reference pulse



Background identification: fit and results

Radioactive contaminations of $^{116}\text{CdWO}_4$ crystal scintillators

Chain	Nuclide	Activity mBq/kg
-------	---------	--------------------

^{232}Th	^{232}Th	0.61(2)
-------------------	-------------------	---------

	^{228}Th	0.022(3)
--	-------------------	----------

^{238}U	^{238}U	0.59(7)
------------------	------------------	---------

	^{234}Th	0.64(7)
--	-------------------	---------

	^{230}Th	0.11(2)
--	-------------------	---------



	^{226}Ra	≤ 0.01
--	-------------------	-------------

	^{210}Pb	0.6(1)
--	-------------------	--------

	^{40}K	0.20(1)
--	-----------------	---------

	$^{110\text{m}}\text{Ag}$	< 0.06
--	---------------------------	----------

Total α activity = 2.27 mBq/kg



Result for two neutrino double beta decay of ^{116}Cd

Conditions of the Fit:

- Variation of bounds for radioactive contaminations
- Model of background
- Interval of fit
- Quenching for β (non prop. light response) [1,2]


[1] PRC 76(2007)064603 [2] NIMA 696 (2012) 144

Signal to bkg ratio: 2.6 in [1.1–2.8] MeV

Systematic errors

Source	SE%
Rad. contamination of $^{116}\text{CdWO}_4$ crystals	65
BG models, MC, QF	15
PSD efficiency	10
Interval of the fit	7
Number of ^{116}Cd nuclei	3

$$T_{1/2} = [2, 63^{+0,11}_{-0,12}] \times 10^{19} \text{ yr} \quad (\text{the most accurate value up to date})$$



ELEGANT: J. Phys. Soc. Japan 64(1995)339
 Solotvina 1995: Phys. Lett. B 344(1995)72
 NEMO-2: Z. Phys. C 72(1996)239
 Solotvina (2000): PRC 62(2000)045501
 Solotvina (2003): PRC 68(2003)035501
 NEMO-2* (recalc.): PRC 81(2010)035501
 NEMO-3: PRD 95(2017)012007

$T_{1/2}$ limit on $0\nu 2\beta$ decay of ^{116}Cd

Further background reduction (**~35%**) for $0\nu 2\beta$ decay by excluding events from:

^{212}Bi [$Q_\alpha=6207.26(3)$ keV, B.R. ~36%] \rightarrow ^{208}Tl [$Q_\beta=4998.9(18)$ keV, $T_{1/2} = 3.053(4)$ min]

\Rightarrow **background rate in 2.7-2.9 MeV: 0.07 (counts/keV/kg/yr)**
(live time reduction ~15%)


Fit in 2.5–3.2 MeV: -3.7 ± 10.6 counts

$T_{1/2} > 2.2 \times 10^{23}$ yr @ 90% C.L.

Effective Majorana neutrino mass:

$\langle m_\nu \rangle < 1.0 - 1.7$ eV [1-4]

+ New improved limits on $T_{1/2}$ for $0\nu 2\beta$ decay to excited levels of ^{116}Sn in the range:
 $(3.6-6.3) \times 10^{22}$ yr



[1] T.R. Rodryguez et al., Phys.Rev.Lett. 105(2010)252503

[2] F. Simkovic et al., Phys.Rev.C 87 (2013)045501

[3] J. Hyvarinen et al., Phys.Rev.C 91 (2015)024613

[4] J. Barea et al., Phys.Rev.C 91(2015)034304

Improvement of radiopurity of $^{116}\text{CdWO}_4$ by recrystallization

A.S. Barabash et al., Nucl. Instr. Meth.A 833(2016)77

Re-crystallized by the low-thermal-gradient Czochralski technique in a platinum crucible



Crystal n.3 used (326 g mass)

60% of initial mass after re-crystallization

Side surface made opaque by grinding paper to improve light collection

Radioactive contamination of the samples (before and after recrystallization) measured in the DAMA/CRYS setup @ LNGS

Chain	Nuclide (sub-chain)	Activity (mBq/kg)	
		Before recrystallization	After recrystallization
^{232}Th	^{232}Th	0.13(7)	0.03(2)
	^{228}Th	0.10(1)	0.010(3)
^{238}U	^{238}U	1.8(2)	0.8(2)
	^{226}Ra	≤ 0.1	≤ 0.015
	$^{234}\text{U} + ^{230}\text{Th}$	0.6(2)	0.4(1)
	^{210}Po	1.6(2)	0.4(1)
Total α		4.44(4)	1.62(4)

➤ ^{228}Th reduced by a factor $\sim 10 \Rightarrow 0.01$ mBq/kg

➤ α activity reduced by a factor $\sim 3 \Rightarrow 1.6$ mBq/kg

main background component for ^{116}Cd
 $0\nu 2\beta$ decay

\Rightarrow Strong segregation of the radioactive elements in the CdWO_4 crystals growing process



THE USED $^{106}\text{CdWO}_4$ CRYSTAL SCINTILLATOR

NIMA615(2010)301

- Samples of cadmium were purified by vacuum distillation (Institute of Physics and Technology, Kharkiv) and the Cadmium tungstate compounds were synthesized from solutions
- Crystal boule was grown by the low-thermal-gradient Czochralski technique (NIIC Novosibirsk) (initial powder 265 g)
- Crystal scintillator (**216 g** mass), **66.4% enrichment in ^{106}Cd** (2.66×10^{23} nuclei of ^{106}Cd) measured by thermal ionisation mass-spectrometry \Rightarrow 2nd enriched CdWO_4 crystal ever produced

1st exp: single crystal in DAMA/R&D


PRC85(2012)044610



2nd exp: coincidence with 4 HP-Ge

PRC93(2016)045502

Stella facility
@LNGS





Search for $\beta\beta$ decay in ^{106}Cd with $^{106}\text{CdWO}_4$ scintillator in coincidence with 4 HPGe (GeMulti)

PRC93 (2016) 045502

^{106}Cd , a promising isotope:

- ✓ One of the six isotopes candidate for $2\beta^+$ decay
- ✓ $\delta = (1.25 \pm 0.06)\%$ \Rightarrow possible enrichment up to 100%
- ✓ $Q_{2\beta} = (2775.39 \pm 0.10)$ keV $\Rightarrow 2\beta^+$, $\epsilon\beta^+$, 2ϵ modes possible
- ✓ Possible resonant $2\epsilon 0\nu$ captures to excited level of ^{106}Pd
- ✓ Theoretical $T_{1/2}$ favorable for some 2ν modes ($10^{20} - 10^{22}$ yr)



$^{106}\text{CdWO}_4$ crystal scintillator:


- ✓ Mass: 216 g, 66.4% enrichment in ^{106}Cd
- ✓ Good scintillation properties
- ✓ Active source approach (high detection efficiency)
- ✓ Low levels of internal contamination in (U, Th K)
- ✓ α/β discrimination capability

PbWO₄ light-guide (Ø40 × 83 mm)


Reduce PMT background (archael. lead: $A(^{210}\text{Pb}) < 0.3$ mBq/kg)

$^{106}\text{CdWO}_4$ crystal scintillator in GeMulti: Results

1. In anticoincidence with the HPGe detectors (**AC**)
2. In coincidence with $E_{\text{HPGe}} > 200$ keV (**CC >200**)
3. In coincidence with $E_{\text{HPGe}} = 511$ keV (**CC 511**)
4. In coincidence with $E_{\text{HPGe}} = 1160$ keV (**CC 1160**)



Energy spectrum of $^{106}\text{CdWO}_4$ detector in coincidence with 511 keV in HPGe (circles). Monte Carlo simulated distributions of 2β decay of ^{106}Cd excluded at 90% CL





- New limits on 2ε , $\varepsilon\beta^+$, $2\beta^+$ processes on the level of $T_{1/2} > 10^{20} - 10^{21}$ yr
- The half-life limit on the $\varepsilon\beta^+2\nu$ decay, $T_{1/2} > 1.1 \times 10^{21}$ yr, reached the region of theoretical predictions
- For $2\varepsilon 0\nu$ resonant captures: $T_{1/2} > (8.5 \times 10^{20} - 1.4 \times 10^{21})$ yr



New $^{106}\text{CdWO}_4$ experiment in DAMA/CRYS set-up

- 1) $^{106}\text{CdWO}_4$ in (anti)coincidence with two large CdWO_4 scintillators mounted in DAMA/CRYS set-up @ LNGS
- 2) High efficiency
- 3) Experiment in data taking since May 2016


DAMA/CRYS set-up



New $^{106}\text{CdWO}_4$ experiment in DAMA/CRYS set-up



ENERGY SPECTRA OF ^{106}Cd




The energy spectra accumulated over **6935 h** by the $^{106}\text{CdWO}_4$ detector:

- in anticoincidence with the $^{nat}\text{CdWO}_4$ detectors
- in coincidence with event(s) in at least one of the $^{nat}\text{CdWO}_4$ detectors with energy:
 - $E > 200$ keV
 - E in energy window around 511 keV
 - E in energy window around 1160 keV


ESTIMATION OF SENSITIVITY

Expected signal for $^{106}\text{Cd } 0\nu 2\beta(0^+ \rightarrow 0^+)$:


Spectrum of $^{106}\text{CdWO}_4$ detector



Spectrum of $^{106}\text{CdWO}_4$ detector when one of the two CdWO_4 detectors detects γ of 511 keV ($\pm 2\sigma$)



Spectrum of $^{106}\text{CdWO}_4$ detector when both the CdWO_4 detectors detect γ of 511 keV ($\pm 2\sigma$)



Sensitivity after 1yr in the hypothesis of about 30 background counts in [0.-3.] MeV:

$0\nu e\beta^+$ (g.s.): $T_{1/2} \approx 5 \times 10^{21}$ yr

$2\nu e\beta^+$ (g.s.): $T_{1/2} \approx 3 \times 10^{21}$ yr



$2\nu 2\beta^+$ (g.s.): $T_{1/2} \approx 2 \times 10^{21}$ yr

In the region of theoretical predictions: $T_{1/2} \sim 10^{20} - 10^{22}$ yr

Note that, up to now, 2ν mode of the $2\beta^+$ processes has not been detected unambiguously: there are only indications for ^{130}Ba and ^{78}Kr


Running and future experiments on HPGE

- Experiment running since February 2015 with deeply purified Nd_2O_3 sample (2381 g) in GeMulti detector to investigate 2β decay of ^{150}Nd to excited levels of ^{150}Sm :
 - ⇒ Background rate in the region of expected peaks (334.0 keV and 406.5 keV) ≈ 2 counts/keV/d
 - ⇒ Expected $T_{1/2}$ sensitivity after 500 days of measurements: 1.3×10^{20} yr (90%CL)



Improvement of Nd_2O_3 radioactive contamination

Contamination	Before [1]	Present
^{40}K	46	< 4
^{214}Bi (^{226}Ra)	1.1	< 0.4
^{228}Ac (^{228}Ra)	0.9	< 0.4




- New experiment to search for 2β of osmium (and α decay of osmium to excited level of daughter nuclei) in progress with BEGe detector:
 - ⇒ Detection efficiency significantly improved by cutting the osmium rods into thin (0.8-1 mm) plates and by using the BEGe detector

- Purification of Er, Yb, and Sm is in progress for experiments to search for resonant $2\varepsilon 0\nu$ processes in these nuclei



2β decay of ^{150}Nd to the excited states of ^{150}Sm



An experiment with highly purified Nd_2O_3 source (2.381 kg) in the GeMulti (4 HPGe ~220 cm^3 each) detector of the STELLA facility



Previous results:

$$T_{1/2}^{2\nu 2\beta} = [1.33^{+0.36}_{-0.23} (\text{st})^{+0.27}_{-0.13} (\text{syst})] \times 10^{20} \text{ yr} \quad [1]$$

$$T_{1/2}^{2\nu 2\beta} = [1.07^{+0.45}_{-0.25} (\text{st}) \pm 0.07 (\text{syst})] \times 10^{20} \text{ yr} \quad [2]$$



8 events of the decay are observed over 16375 in coincidence with the HPGe. The half-life:

$$T_{1/2}^{2\nu 2\beta} = (0.84^{+0.47}_{-0.22}) \times 10^{20} \text{ yr}$$

The experiment in progress aiming to improve the statistic accuracy to at least 3σ


[1] PRC 79 (2009) 045501

[2] PRC 90 (2014) 055501

Running and future experiments


ZnWO_4 running in DAMA/R&D

- New 4 crystal scintillators in the DAMA/R&D in data taking:
 - radioactive contamination and scintillation performances
 - study of double beta decay modes in Zn and W isotopes



Development of Detectors with Anisotropic Response for Dark Matter Search in Directionality Approach are in progress

$^{106}\text{CdWO}_4$ running in DAMA/CRYS



$^{106}\text{CdWO}_4$ in (anti)coincidence with two large CdWO_4 scintillators to study $\beta\beta$ decays of ^{106}Cd



Thanks for your attention